Index of Authors

ABRAHAM, E. P. *see* HUDDLESTON, J. A. 705–707
ADELMANN, B. C. *see* DESSAU, W. 55–59
AKHTAR, M., ALEXANDER, K., BOAR, R. B., MCGHIE, J. F. & BARTON, D. H. R. Chemical and enzymic studies on the characterization of intermediates during the removal of the 14α-methyl group in cholesterol biosynthesis. The use of 32-functionalized lanostane derivatives 449–463
ALEXANDER, K. *see* AKHTAR, M. 449–463
ALI, S. Y. *see* BAYLISS, M. T. 123–132
ANDERSON, P. J. Actin in young and senescent fibroblasts 169–172
ANTONINI, E. *see* SILVESTRINI, M. C. 257–263
ARCHIBALD, A. R. *see* HEPTINSTALL, J. 329–336
AUDELL, L. *see* INEROT, S. 143–156
AURICCHIO, F., ROTONDI, A., SCHIAVONE, E. & BRESCIANI, F. Oestrogen receptor of mammary gland. Inhibition of aggregation and characterization of receptor from lactating gland in the presence of sodium bromide 481–488
AXELSEN, N. H. *see* FOLTMANN, B. 425–427
AXELSSON, I. & HEINEGÅRD, D. Characterization of the keratan sulphate proteoglycans from bovine corneal stroma 517–530
AYALA, J. A. & NIETO, M. Thermal denaturation of *Micrococcus lysodeikticus* adenosine triphosphatase. Influence of temperature on the circular dichroism, fluorescence and enzymic activity of the protein 371–380

BADDILEY, J. *see* HEPTINSTALL, J. 329–336
BAISTED, D. J. *see* STAVER, M. J. 297–303
BALDWIN, S. A. & PERHAM, R. N. Novel kinetic and structural properties of the class-I D-fructose 1,6-bisphosphate aldolase from *Escherichia coli* (Crookes' strain) 643–652
BALDWIN, S. A., PERHAM, R. N. & STRIBLING, D. Purification and characterization of the class-II D-fructose 1,6-bisphosphate aldolase from *Escherichia coli* (Crookes' Strain) 633–641
BAMFORTH, C. W. & QUAYLE, J. R. The dye-linked alcohol dehydrogenase of *Rhodopseudomonas acidophila*. Comparison with dye-linked methanol dehydrogenases 677–686
BARCLAY, A. N. *see* KUCHEL, P. W. 411–417
BARRETT, A. J. *see* ROUGHLEY, P. J. 721–724
BARTHOLOMEW, B., DODGSON, K. S. & GORHAM, S. D. Purification and properties of the S1 secondary alkylsulphohydrolase of the detergent-degrading microorganism, *Pseudomonas* C12B 659–667
BARTON, D. H. R. *see* AKHTAR, M. 449–463
BAUERMEISTER, A. *see* NIMMO, I. A. 437–440
BAYLISS, M. T. & ALI, S. Y. Isolation of proteoglycans from human articular cartilage 123–132
BEINERT, H. *see* STEENKAMP, D. J. 361–369
BERGSMA, D. R. *see* WIGGERT, B. 87–94
BIELER, L. Z. *see* CAMMER, W. 567–577
BIRDEN, R. *see* BRISTOW, A. F. 381–388
BOAR, R. B. *see* AKHTAR, M. 449–463
BOER, P. *see* REUVERS, F. 505–508
BONORI, M. *see* RIGO, A. 277–280
BRAND, R. C., KLOOTWIJK, J., PLANTA, R. J. & MADEN, B. E. H. Biosynthesis of a hypermodified nucleotide in *Saccharomyces carlsbergensis* 17S and HeLa-cell 18S ribosomal ribonucleic acid 71–77
BRAY, R. C. *see* LOWE, D. J. 471–479
BRESCIANI, F. *see* AURICCHIO, F. 481–488
BREWER, G. J. *see* GILMAN, J. G. 625–632
BRISTOW, A. F. & VIRDEN, R. Preferential nitration with tetranitromethane of a specific tyrosine residue in penicillinase from *Staphylococcus aureus* PC1. Evidence that the preferentially nitrated residue is not part of the active site but that loss of activity is due to intermolecular cross-linking 381–388
BRITTON, H. G. & DANN, L. G. Use of flux ratio measurements for the determination of the order of addition of substrates and products in enzyme reactions 29–37
BRITTON, H. G. *see also* DANN, L. G. 39–54
BRUNORI, M. *see* SILVESTRINI, M. C. 257–263

CAKE, M. H. *see* LITWACK, G. 445–448
CALABRESE, L. *see* RIGO, A. 277–280
CAMMER, W., BIELER, L. Z. & NORTON, W. T. Proteolytic and peroxidatic reactions of commercial horseradish peroxidase with myelin basic protein 567–577
CAMPBELL, D. G. *see* KUCHEL, P. W. 411–417
CAPPUCCINELLI, P. & HAMES, B. D. Characterization of colchicine-binding activity in *Dictyostelium discoideum* 499–504
CHADER, G. J. *see* WIGGERT, B. 87–94
CHANDLER, H. D., WOOLF, C. J. & HEPBURN, H. R. Gliding edge dislocations in proteins as a mechanism for active ion transport 559–565
COCCO, D. *see* RIGO, A. 277–280
COGGINS, J. R. *see* LUMSDEN, J. 441–444
COLE, R. D. *see* HODSDON, J. 429–432
COLEY, J. *see* HEPTINSTALL, J. 329–336
COLOSIMO, A. *see* SILVESTRINI, M. C. 257–263

DANIELS, M. *see* WALLIS, M. 669–676
DANN, L. G. & BRITTON, H. G. Kinetics and mechanism of action of muscle pyruvate kinase 39–54
DANN, L. G. *see also* BRITTON, H. G. 29–37
DAVIES, R. V. *see* WALLIS, M. 669–676
DE JONG, P. J. *see* KLAUSNER, Y. S. 157–167
DEL VILLAR, E. *see* SANCHEZ, E. 173–177
DESSAU, W., ADELMANN, B. C., TIMPL, R. & MARTIN, G. R. Identification of the sites in collagen α-chains that bind serum anti-gelatin factor (cold-insoluble globulin) 55–59
DIXON, H. B. F. *see* WEBSTER, D. 239–244
DODGSON, K. S. *see* BARTHOLOMEW, B. 659–667
DOWER, S. K., GETTINS, P., JACKSON, R., DWEK, R. A. & GIVOL, D. The binding of 2,4,6-trinitrophenyl derivatives to the mouse myeloma immunoglobulin A protein MOPC 315 179–188
DWEK, R. A. *see* DOWER, S. K. 179–188

EASTERBROOK-SMITH, S. B., WALLACE, J. C. & KEECH, D. B. A reappraisal of the reaction pathway of pyruvate carboxylase 225–228

ELDER, G. H. & EVANS, J. O. A radiochemical method for the measurement of coproporphyrinogen oxidase and the utilization of substrates other than coproporphyrinogen III by the enzyme from rat liver 205–214
ELDER, G. H., EVANS, J. O., JACKSON, J. R. & JACKSON, A. H. Factors determining the sequence of oxidative decarboxylation of the 2- and 4-propionate substituents of coproporphyrinogen III by coproporphyrinogen oxidase in rat liver 215–223
EL-HAMALAWI, A.-R. A. Isolation and separation of α- and β-tubulin from chick-embryo brain, muscle and skin 717–719
EVANS, J. O. *see* ELDER, G. H. 205–214, 215–223
EVANSON, J. M. *see* WOOLLEY, D. E. 265–276

FILLER, R. *see* LITWACK, G. 445–448
FOLTMANN, B., LØNBLAD, P. & AXELSEN, N. H. Demonstration of chymosin (EC 3.4.23.4) in the stomach of newborn pig 425–427
FREYSSINET, J.-M., LEWIS, B. A., HOLBROOK, J. J. & SHORE, J. D. Protein–protein interactions in blood clotting. The use of polarization of fluorescence to measure the dissociation of plasma factor XIIIa 403–410
FREYSSINET, J.-M. *see also* LEWIS, B. A. 397–402

GEBHARDT, O. *see* WALLIN, R. 95–101
GETTINS, P. *see* DOWER, S. K. 179–188
GILMAN, J. G. & BREWER, G. J. The oxygen-linked zinc-binding site of human haemoglobin 625–632
GIVOL, D. *see* DOWER, S. K. 179–188
GLANVILLE, R. W. *see* WOOLLEY, D. E. 265–276
GLICK, K. *see* STAVER, M. J. 297–303
GOODMAN, S. L., ISERN DE CALDENTEY, M. & WHEELER, K. P. A simple and rapid method for the reversible removal of lipids from a membrane-bound enzyme 305–311
GOODMAN, S. L. & WHEELER, K. P. Ouabain binding to phospholipid-dependent adenosine triphosphatase 313–320
GORHAM, S. D. *see* BARTHOLOMEW, B. 659–667
GRAHAM, M. *see* KAPLOWITZ, N. 465–470
GREENBERG, R. *see* GROVES, M. L. 337–342
GRIFFIN, M. *see* HALLIWELL, G. 713–715
GROVES, M. L. & GREENBERG, R. Isolation, composition and ordering of cyanogen bromide peptides in human casein 337–342

HALLIWELL, G. & GRIFFIN, M. Affinity chromatography of the cellulase system of *Trichoderma koningii* 713–715
HAMES, B. D. *see* CAPPUCCINELLI, P. 499–504
HATTON, M. W. C. *see* KOJ, A. 589–596
HEINEGÅRD, D. *see* AXELSSON, I. 517–530; INEROT, S. 143–156
HELMSEN, R. *see* WIGGERT, B. 87–94
HEMMING, F. W. *see* REUVERS, F. 505–508
HEPBURN, H. R. *see* CHANDLER, H. D. 559–565
HEPTINSTALL, J., COLEY, J., WARD, P. J., ARCHIBALD, A. R. & BADDILEY, J. The linkage of sugar phosphate polymer to peptidoglycan in walls of *Micrococcus* sp. 2102 329–336
HIBASAMI, H. & PEGG, A. E. Rapid and convenient method for the assay of aminopropyltransferases 709–712
HIRSCH, J. & MARTELO, O. J. Purification and properties of a nuclear protein kinase associated with ribonucleic acid polymerase I 355–359
HODSDON, J., KOLB, H., SNELL, E. E. & COLE, R. D. The pyridoxal-binding site in pyridoxamine–pyruvate transaminase 429–432
HOLBROOK, J. J. *see* FREYSSINET, J.-M. 403–410; LEWIS, B. A. 397–402; LODOLA, A. 577–588
HUDDLESTON, J. A., ABRAHAM, E. P., YOUNG, D. W., MORECOMBE, D. J. & SEN, P. K. The stereochemistry of β-lactam formation in cephalosporin biosynthesis 705–707
HURST, P. L., SULLIVAN, P. A. & SHEPHERD, M. G. Substrate specificity and mode of action of a cellulase from *Aspergillus niger* 389–395
HUTSON, N. J. *see* SUGDEN, P. H. 433–435

INEROT, S., HEINEGÅRD, D., AUDELL, L. & OLSSON, S.-E. Articular-cartilage proteoglycans in aging and osteoarthritis 143–156
ISERN DE CALDENTEY, M. *see* GOODMAN, S. L. 305–311
IYER, K. S., LAU, S.-J., LAURIE, S. H. & SARKAR, B. Synthesis of the native copper(II)-transport site of human serum albumin and its copper(II)-binding properties 61–69

JACKSON, A. H. *see* ELDER, G. H. 215–223
JACKSON, J. R. *see* ELDER, G. H. 215–223
JACKSON, R. *see* DOWER, S. K. 179–188
JARDINE, N. J. & LEAVER, J. L. The fractionation of histones isolated from *Euglena gracilis* 103–111
JASANI, B., JASANI, M. K. & TALBOT, M. D. Characterization of two acid proteinases found in rabbit skin homografts 287–295
JASANI, M. K. *see* JASANI, B. 287–295
JONES, A. J. S. & RUMSBY, M. G. Interaction of the myelin basic protein with the anionic detergent sodium dodecyl sulphate 281–285
JONES, I. G. *see* LONSDALE, D. M. 79–85
JØRGENSEN, K. E., LIND, K. E., RØIGAARD-PETERSEN, H. & MØLLER, J. V. The functional unit of calcium-plus-magnesium-ion-dependent adenosine triphosphatase from sarcoplasmic reticulum. The aggregational state of the deoxycholate-solubilized protein in an enzymically active form 489–498

KAPLOWITZ, N., SPINA, C., GRAHAM, M. & KUHLENKAMP, J. Glutathione *S*-transferase in human lymphoid cell lines and fractionated peripheral leucocytes 465–470
KEECH, D. B. *see* EASTERBROOK-SMITH, S. B. 225–228
KERBEY, A. L. *see* SUGDEN, P. H. 433–435
KETTERER, B. *see* TIPPING, E. 509–516
KÈZDY, F. J. *see* PANAYOTATOS, N. 421–423
KHAN, M. S. N., SALIM, M. & MADEN, B. E. H. Extensive homologies between the methylated nucleotide sequences in several vertebrate ribosomal ribonucleic acids 531–542
KIDO, R. *see* NOGUCHI, T. 113–122
KIENER, P. A. & WALEY, S. G. Reversible inhibitors of penicillinases 197–204
KLAUSNER, Y. S., RIGBI, M., TICHO, T., DE JONG, P. J., NEGINSKY, E. J. & RINOTT, Y. The interaction of α-*N*-(*p*-toluenesulphonyl)-*p*-guanidino-L-phenylalanine methyl ester with thrombin and trypsin 157–167

KLOOTWIJK, J. *see* BRAND, R. C. 71–77
KNOWLES, P. F. *see* VIRJI, M. 343–353
KOJ, A., HATTON, M. W. C., WONG, K.-L. & REGOECZI, E. Isolation and partial characterization of rabbit plasma α_1-antitrypsin 589–596
KOLB, H. *see* HODSDON, J. 429–432
KOSKELO, P. *see* TIPPING, E. 509–516
KRAUS, J. Recognition of individual *Escherichia coli* transfer ribonucleic acids by 1-adenine-specific methyltransferase from rat liver 247–249
KUCHEL, P. W., CAMPBELL, D. G., BARCLAY, A. N. & WILLIAMS, A. F. Molecular weights of the Thy-1 glycoproteins from rat thymus and brain in the presence and absence of deoxycholate 411–417
KUHLENKAMP, J. *see* KAPLOWITZ, N. 465–470
KYFFIN, T. W. *see* SCOTT, J. E. 697–701

LAMPERT, J. *see* LANDMAN, A. D. 255–256
LANDMAN, A. D. & LAMPERT, J. Activation of immobilized acetyl-coenzyme A carboxylase 255–256
LAU, S.-J. *see* IYER, K. S. 61–69
LAURIE, S. H. *see* IYER, K. S. 61–69
LEAVER, J. L. *see* JARDINE, N. J. 103–111
LEHTOVAARA, P. Anomalous migration of leghaemoglobin on sodium dodecyl sulphate/polyacrylamide-gel electrophoresis 251–253
LEWIS, B. A., FREYSSINET, J.-M. & HOLBROOK, J. J. An equilibrium study of metal ion binding to human plasma coagulation Factor XIII 397–402
LEWIS, B. A. *see also* FREYSSINET, J.-M. 403–410
LIBOR, S. M., SUNDARAM, T. K. & SCRUTTON, M. C. Pyruvate carboxylase from a thermophilic *Bacillus*. Studies on the specificity of activation by acyl derivatives of coenzyme A and on the properties of catalysis in the absence of activator 543–558
LIND, K. E. *see* JØRGENSEN, K. E. 489–498
LITWACK, G., CAKE, M. H., FILLER, R. & TAYLOR, K. Physical measurements of the liver glucocorticoid receptor 445–448
LODOLA, A., SPRAGG, S. P. & HOLBROOK, J. J. Malate dehydrogenase of the cytosol. Preparation and reduced nicotinamide–adenine dinucleotide-binding studies 577–588
LØNBLAD, P. *see* FOLTMANN, B. 425–427
LONSDALE, D. M. & JONES, I. G. Localization of the ribonucleotide sites in rat liver mitochondrial deoxyribonucleic acid 69–85
LOWE, D. J. & BRAY, R. C. Magnetic coupling of the molybdenum and iron–sulphur centres in xanthine oxidase and xanthine dehydrogenases 471–479
LUMSDEN, J. & COGGINS, J. R. The subunit structure of the *arom* multienzyme complex of *Neurospora crassa*. Evidence from peptide 'maps' for the identity of the subunits 441–444
LYNEN, A., SEDLACZEK, E. & WIELAND, O. H. Partial purification and characterization of a pyruvate dehydrogenase-complex-inactivating enzyme from rat liver 321–328

MADEN, B. E. H. *see* BRAND, R. C. 71–77; KHAN, M. S. N. 531–542
MARTELO, O. J. *see* HIRSCH, J. 355–359
MARTIN, G. R. *see* DESSAU, W. 55–59
MCGHIE, J. F. *see* AKHTAR, M, 449–463
MCTAGGART, F. *see* WEECH, P. K. 687–695
MILLS, G. L. *see* WEECH, P. K. 687–695
MINATOGAWA, Y. *see* NOGUCHI, T. 113–122
MØLLER, J. V. *see* JØRGENSEN, K. E. 489–498
MORECOMBE, D. J. *see* HUDDLESTON, J. A. 705–707
MURPHY, G. *see* ROUGHLEY, P. J. 721–724

NEGINSKY, E. J. *see* KLAUSNER, Y. S. 157–167
NIETO, M. *see* AYALA, J. A. 371–380
NIEUWENHUIZEN, W. *see* VAN RUIJVEN-VERMEER, I. A. M. 653–658
NIMMO, I. A. & BAUERMEISTER, A. A theoretical analysis of the use of zonal gel filtration in the detection and purification of protein–ligand complexes 437–440
NOGUCHI, T., OKUNO, E., TAKADA, Y., MINATOGAWA, Y., OKAI, K. & KIDO, R. Characteristics of hepatic alanine–glyoxylate aminotransferase in different mammalian species 113–122
NORTON, W. T. *see* CAMMER, W. 567–577

OKAI, K. *see* NOGUCHI, T. 113–122
OKUNO, E. *see* NOGUCHI, T. 113–122
OLSSON, S.-E. *see* INEROT, S. 143–156

PANAYOTATOS, N. & KÈZDY, F. J. Kinetics and template-dependency of ribonucleic acid synthesis by bacterial ribonucleic acid polymerase 421–423
PEGG, A. E. *see* HIBASAMI, H. 709–712
PERHAM, R. N. *see* BALDWIN, S. A. 633–641; 643–652
PLANTA, R. J. *see* BRAND, R. C. 71–77
PRYDZ, H. *see* WALLIN, R. 95–101

QUAYLE, J. R. *see* BAMFORTH, C. W. 677–686

RANDLE, P. J. *see* SUGDEN, P. H. 433–435
REGOECZI, E. *see* KOJ, A. 589–596
REUVERS, F., BOER, P. & HEMMING, F. W. The presence of dolichol in a lipid diphosphate *N*-acetylglucosamine from *Saccharomyces cerevisiae* (Baker's Yeast) 505–508
RIGBI, M. *see* KLAUSNER, Y. S. 157–167
RIGO, A., VIGLINO, P., BONORI, M., COCCO, D., CALABRESE, L. & ROTILIO, G. The binding of copper ions to copper-free bovine superoxide dismutase. Kinetic aspects 277–280
RINOTT, Y. *see* KLAUSNER, Y. S. 157–167
RISTELI, L. Further characterization of collagen galactosyltransferase from chick embryos 189–196
ROBERTS, D. R. *see* WOOLLEY, D. E. 265–276
ROBERTS, N. B. & TAYLOR, W. H. The isolation and properties of a non-pepsin proteinase from human gastric mucosa 617–624
ROBERTS, N. B. & TAYLOR, W. H. The preparation and purification of individual human pepsins by using diethylaminoethyl-cellulose 607–615
RØIGAARD-PETERSEN, H. *see* JØRGENSEN, K. E. 489–498
ROTILIO, G. *see* RIGO, A. 277–280
ROTONDI, A. *see* AURICCHIO, F. 481–488
ROUGHLEY, P. J., MURPHY, G. & BARRETT, A. J. Proteinase inhibitors of bovine nasal cartilage 721–724
RUMSBY, M. G. *see* JONES, A. J. S. 281–285

SALIM, M. *see* KHAN, M. S. N. 531–542

SANCHEZ, E., DEL VILLAR, E. & TEPHLY, T. R. Structural requirements in the reaction of morphine uridine diphosphate glucuronyltransferase with opioid substances 173–177
SARKAR, B. *see* IYER, K. S. 61–69
SCHIAVONE, E. *see* AURICCHIO, F. 481–488
SCOTT, J. E. & KYFFIN, T. W. Demineralization in organic solvents by alkylammonium salts of ethylenediaminetetra-acetic acid 697–701
SCRUTTON, M. C. *see* LIBOR, S. M. 543–558
SEDLACZEK, E. *see* LYNEN, A. 321–328
SEN, P. K. *see* HUDDLESTON, J. A. 705–707
SHEPHERD, M. G. *see* HURST, P. L. 389–395
SHORE, J. D. *see* FREYSSINET, J.-M. 403–410
SILVESTRINI, M. C., COLOSIMO, A., BRUNORI, M. & ANTONINI, E. Properties of cytochrome *c* modified by attachment to a carbohydrate polymer 257–263
SINGER, T. P. *see* STEENKAMP, D. J. 361–369
SNELL, E. E. *see* HODSDON, J. 429–432
SPARKES, M. J. *see* WEBSTER, D. 239–244
SPINA, C. *see* KAPLOWITZ, N. 465–470
SPRAGG, S. P. *see* LODOLA, A. 577–588
STANSFIELD, D. A. *see* YOUNG, J. L. 133–142
STAVER, M. J., GLICK, K. & BAISTED, D. J. Uridine diphosphate glucose–sterol glucosyltransferase and nucleoside diphosphatase activities in etiolated pea seedlings 297–303
STEENKAMP, D. J., SINGER, T. P. & BEINERT, H. Participation of the iron–sulphur cluster and of the covalently bound coenzyme of trimethylamine dehydrogenase in catalysis 361–369
STRIBLING, D. *see* BALDWIN, S. A. 633–641
SUGDEN, P. H., HUTSON, N. J., KERBEY, A. L. & RANDLE, P. J. Phosphorylation of additional sites on pyruvate dehydrogenase inhibits its re-activation by pyruvate dehydrogenase phosphate phosphatase 433–435
SULLIVAN, P. A. *see* HURST, P. L. 389–395
SUNDARAM, T. K. *see* LIBOR, S. M. 543–558

TAKADA, Y. *see* NOGUCHI, T. 113–122
TALBOT, M. D. *see* JASANI, B. 287–295
TAYLOR, K. *see* LITWACK, G. 445–448
TAYLOR, W. H. *see* ROBERTS, N. B. 607–615; 617–624
TEPHLY, T. R. *see* SANCHEZ, E. 173–177
TICHO, T. *see* KLAUSNER, Y. S. 157–167
TIMPL, R. *see* DESSAU, W. 55–59
TIPPING, E., KETTERER, B. & KOSKELO, P. The binding of porphyrins by ligandin 509–516
TJERNSHAUGEN, H. Partial purification and properties of an acid nucleotidase from the postmicrosomal supernatant of rat spleen 597–605

VAN RUIJVEN-VERMEER, I. A. & NIEUWENHUIZEN, W. Purification of rat fibrinogen and its constituent chains 653–658
VIGLINO, P. *see* RIGO, A. 277–280
VIRJI, M. & KNOWLES, P. F. Protein–lipid interactions in cytochrome oxidase from *Saccharomyces cerevisiae*. Effects of detergents and reconstitution of enzyme activity by phospholipids by using cholate-mediated exchange 343–353

WALEY, S. G. *see* KIENER, P. A. 197–204
WALLACE, J. C. *see* EASTERBROOK-SMITH, S. B. 225–228
WALLIN, R., GEBHARDT, O. & PRYDZ, H. NAD(P)H dehydrogenase and its role in the vitamin K (2-methyl-3-phytyl-1,4-naphthaquinone)-dependent carboxylation reaction 95–101
WALLIS, M., DAVIES, R. V. & DANIELS, M. Preparation of internally labelled rat pituitary somatotropin (growth hormone) 669–676
WARD, P. J. *see* HEPTINSTALL, J. 329–336
WEBSTER, D., SPARKES, M. J. & DIXON, H. B. F. An arsenical analogue of adenosine diphosphate 239–244
WEECH, P. K., MCTAGGART, F. & MILLS, G. L. Comparison of guinea-pig serum lipoproteins after iodination by two different methods 687–695
WHEELER, K. P. *see* GOODMAN, S. L. 305–311, 313–320
WIELAND, O. H. *see* LYNEN, A. 321–328
WIGGERT, B., BERGSMA, D. R., HELMSEN, R. & CHADER, G. J. Vitamin A receptors. Retinoic acid binding in ocular tissues 87–94
WILKINSON, J. M. The components of troponin from chicken fast skeletal muscle. A comparison of troponin T and troponin I from breast and leg muscle 229–238
WILLIAMS, A. F. *see* KUCHEL, P. W. 411–417
WONG, K.-L. *see* KOJ, A. 589–596
WOOLF, C. J. *see* CHANDLER, H. D. 559–565
WOOLLEY, D. E., GLANVILLE, R. W., ROBERTS, D. R. & EVANSON, J. M. Purification, characterization and inhibition of human skin collagenase 265–276

YOUNG, D. W. *see* HUDDLESTON, J. A. 705–707
YOUNG, J. L. & STANSFIELD, D. A. Stabilization and solubilization of bovine corpus-luteum adenylate cyclase. The effects of guanosine triphosphate, guanosine 5′-[β,γ-imido]triphosphate, sodium fluoride and Tris/hydrochloric acid concentration on enzyme activity 133–142

Index of Subjects

Acetyl-coenzyme A carboxylase, adipose-tissue, rat, immobilized, activation of (Landman, A. D. & Lampert, J.) 255–256

Acid nucleotidase, *see* Nucleotidase, acid

Actin, content of, in young and senescent chick-embryo fibroblasts (Anderson, P. J.) 169–172

Acyl-coenzyme A, specificity of activation by, of thermophilic *Bacillus* sp. pyruvate carboxylase and properties of catalysis by the enzyme in the absence of activator (Libor, S. M., Sundaram, T. K. & Scrutton, M. C.) 543–558

Adenosine diphosphate, synthesis of an arsenical analogue of (Webster, D., Sparkes, M. J. & Dixon, H. B. F.) 239–244

Adenosine triphosphatase, calcium-plus-magnesium ion-dependent, sarcoplasmic-reticulum, skeletal-muscle, rabbit, deoxycholate-solubilized, aggregational state of, in an enzymically active form (Jørgensen, K. E., Lind, K. E., Røigaard-Petersen, H. & Møller, J. V.) 489–498

Adenosine triphosphatase, *Micrococcus lysodeikticus* N.C.T.C. 2665, effect of temperature on the circular dichroism, fluorescence and enzymic activity of (Ayala, J. A. & Nieto, M.) 371–380

Adenosine triphosphatase, sodium-plus-potassium ion-dependent, membrane, kidney, rabbit, simple and rapid method for the reversible removal of lipids from (Goodman, S. L., Isern de Caldentey, M. & Wheeler, K. P.) 305–311

Adenosine triphosphatase, sodium-plus-potassium ion-dependent, phospholipid-dependent, membrane, kidney, rabbit, binding of ouabain to (Goodman, S. L. & Wheeler, K. P.) 313–320

Adenylate cyclase, corpus-luteum, ovary, cow, effects of guanosine triphosphate and other compounds on the activity of (Young, J. L. & Stansfield, D. A.) 133–142

Adipose tissue, rat, activation of immobilized acetyl-coenzyme A carboxylase from (Landman, A. D. & Lampert, J.) 255–256

Aging, changes in the composition of dog articular-cartilage proteoglycans during osteoarthritis and (Inerot, S., Heinegård, D., Audell, L. & Olsson, S.-E.) 143–156

Alanine–glyoxylate aminotransferase, mitochondrial, liver, dog, cat rat and mouse, characteristics of isoenzymes of (Noguchi, T., Okuno, E., Takada, Y., Minatogawa, Y., Okai, K. & Kido, R.) 113–122

Albumin, serum, human, synthesis of and binding of copper ions by a tripeptide corresponding to the copper(II)-binding site of (Iyer, K. S., Lau, S.-J., Laurie, S. H. & Sarkar, B.) 61–69

Alcohol dehydrogenase, dye-linked, *Rhodopseudomonas acidophila*, partial purification of, and comparison of its properties with those of dye-linked methanol dehydrogenases from other micro-organisms (Bamforth, C. W. & Quayle, J. R.) 677–686

Aldolase, *see* Fructose 1,6-bisphosphate aldolase

Alkylammonium ethylenediaminetetra-acetates, demineralization of bone in organic solvents by (Scott, J. E. & Kyffin, T. W.) 697–701

Alkylsulphatase, secondary, S1, *Pseudomonas* C12B, purification and properties of (Bartholomew, B., Dodgson, K. S. & Gorham, S. D.) 659–667

Amino acid sequence, determination of, around the pyridoxal-binding site of *Pseudomonas* MA-1 pyridoxamine–pyruvate aminotransferase (Hodsdon, J., Kolb, H., Snell, E. E. & Cole, R. D.) 429–432

Aminopropyltransferases, rapid method for the assay of the activity of, from various rat tissues (Hibasami, H. & Pegg, A. E.) 709–712

Anti-gelatin factor (cold-insoluble globulin), serum, guinea-pig and human, identification of the sites in α-chains of collagens from various sources that bind (Dessau, W., Adelmann, B. C., Timpl, R. & Martin, G. R.) 55–59

α_1-Anti-trypsin, plasma, rabbit, isolation and partial characterization of (Koj, A., Hatton, M. W. C., Wong, K.-L. & Regoeczi, E.) 589–596

arom multienzyme complex, *Neurospora crassa*, evidence from peptide 'maps' for the identity of subunits of (Lumsden, J. & Coggins, J. R.) 441–444

Arsenical analogue of adenosine diphosphate, synthesis of (Webster, D., Sparkes, M. J. & Dixon, H. B. F.) 239–244

Aspergillus niger, substrate specificity and mode of action of a cellulase from (Hurst, P. L., Sullivan, P. A. & Shepherd, M. G.) 389–395

Bacillus cereus, reversible inhibitors of β-lactamase I from (Kiener, P. A. & Waley, S. G.) 197–204

Bacillus sp., thermophilic, specificity of activation of pyruvate carboxylase from, by acyl-coenzyme A and properties of catalysis by the enzyme in the absence of activator (Libor, S. M., Sundaram, T. K. & Scrutton, M. C.) 543–558

Bacterium W3A1, participation of the iron–sulphur centres and of covalently bound flavin mononucleotide in the reaction catalysed by trimethylamine dehydrogenase from (Steenkamp, D. J., Singer, T. P. & Beinert, H.) 361–369

Basic protein, *see* Protein, basic

Blood plasma, human, equilibrium study of the binding of metal ions to Factor XIII from (Lewis, B. A., Freyssinet, J.-M. & Holbrook, J. J.) 397–402

Blood plasma, human, use of polarization of fluorescence to measure the dissociation of Factor XIIIa from (Freyssinet, J.-M., Lewis, B. A., Holbrook, J. J. & Shore, J. D.) 403–410

Blood plasma, rabbit, isolation and partial characterization of α_1-anti-trypsin from (Koj, A., Hatton, M. W. C., Wong, K.-L. & Regoeczi, E.) 589–596

Blood plasma, rat, purification of fibrinogen and its constituent chains from (van Ruijven-Vermeer, I. A. M. & Nieuwenhuizen, W.) 653–658

Blood serum, guinea-pig and human, identification of the sites in α-chains of collagens from various sources that bind anti-gelatin factor (cold-insoluble globulin) from (Dessau, W., Adelmann, B. C., Timpl, R. & Martin, G. R.) 55–59

Bone, demineralization of, in organic solvents by alkylammonium salts of ethylenediaminetetra-acetic acid (Scott, J. E. & Kyffin, T. W.) 697–701

Brain, chick-embryo, isolation and separation of α- and β-tubulins from, and other tissues (El-Hamalawi, A.-R. A.) 717–719

Brain, ox, cat, rabbit and rat, proteolytic and peroxidatic reactions of commercial preparations of horseradish peroxidase with myelin basic protein from (Cammer, W., Bieler, L. Z. & Norton, W. T.) 567–577

Brain, ox, interaction of sodium dodecyl sulphate with myelin basic protein from (Jones, A. J. S. & Rumsby, M. G.) 281–285

Brain, rat, molecular weights of the Thy-1 membrane glycoproteins from, and thymus gland measured in the presence and absence of deoxycholate (Kuchel, P. W., Campbell, D. G., Barclay, A. N. & Williams, A. F.) 411–417

Brain, rat, rapid method for the assay of the activity of aminopropyltransferases from, and other tissues (Hibasami, H. & Pegg, A. E.) 709–712

Calcium ions, equilibrium study of the binding of, and other metal ions to human plasma Factor XIII (Lewis, B. A., Freyssinet, J.-M. & Holbrook, J. J.) 397–402

Cartilage, articular, dog, changes in the composition of proteoglycans from, during aging and osteoarthritis (Inerot, S., Heinegård, D., Audell, L. & Olsson, S.-E.) 143–156

Cartilage, articular, human, isolation of proteoglycans from (Bayliss, M. T. & Ali, S. Y.) 123–132

Cartilage, nasal-septum, ox, characterization of proteinase inhibitors of (Roughley, P. J., Murphy, G. & Barrett, A. J.) 721–724

Casein, human, isolation, composition and ordering of the peptides obtained by cyanogen bromide cleavage of (Groves, M. L. & Greenberg, R.) 337–342

Cathepsin E, gastric-mucosa, human, isolation and properties of (Roberts, N. B. & Taylor, W. H.) 617–624

Cathepsins, skin, rabbit, two, characterization of (Jasani, B., Jasani, M. K. & Talbot, M. D.) 287–295

Cell wall, *Micrococcus* sp. 2102, linkage of sugar phosphate polymer to peptidogylcan in (Heptinstall, J., Coley, J., Ward, P. J., Archibald, A. R. & Baddiley, J.) 329–336

Cells, HeLa, biosynthesis of a hypermodified nucleotide residue in *Saccharomyces cerevisiae* 17S ribosomal ribonucleic acid and in 18S ribonucleic acid of (Brand, R. C., Klootwijk, J., Planta, R. J. & Maden, B. E. H.) 71–77

Cells, lymphoid, human, activities of glutathione *S*-transferases in fractionated human peripheral leucocytes and (Kaplowitz, N., Spina, C., Graham, M. & Kuhlenkamp, J.) 465–470

Cells, vertebrate, various, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542

Cells, *see also* Fibroblasts

Cellulase, *Aspergillus niger*, substrate specificity and mode of action of (Hurst, P. L., Sullivan, P. A. & Shepherd, M. G.) 389–395

Cellulase system, *Trichoderma koningii*, affinity chromatography of the components of (Halliwell, G. & Griffin, M.) 713–715

Cephalosporium acremonium, stereochemistry of the formation of the β-lactam ring during the biosynthesis of cephalosporin by (Huddleston, J. A., Abraham, E. P., Young, D. W., Morecombe, D. J. & Sen, P. K.) 705–707

Cephalosporium, stereochemistry of the formation of the β-lactam ring during the biosynthesis of, by *Cephalosporium acremonium* (Huddleston, J. A., Abraham, E. P., Young, D. W., Morecombe, D. J. & Sen, P. K.) 705–707

Chick embryo, characterization of collagen galactosyltransferase from (Risteli, L.) 189–196

Chick-embryo fibroblasts, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from, and other vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542

Chick-embryo fibroblasts, young and senescent, content of actin in (Anderson, P. J.) 169–172

Chick embryo, isolation and separation of α- and β-tubulins from brain, muscle and skin from (El-Hamalawi, A.-R. A.) 717–719

Chicken, comparison of troponin T and troponin I from breast and leg skeletal muscle from (Wilkinson, J. M.) 229–238

Cholesterol, use of 32-functionalized derivatives of lanostane in chemical and enzymic studies on the characterization of intermediates during the removal of the 14α-methyl group in the biosynthesis of, by rat liver cell-free preparations (Akhtar, M., Alexander, K., Boar, R. B., McGhie, J. F. & Barton, D. H. R.) 449–463

Chymosin, *see* Rennin

Clotting Factor XIII, plasma, human, equilibrium study of the binding of metal ions to (Lewis, B. A., Freyssinet, J.-M. & Holbrook, J. J.) 397–402

Clotting Factor XIIIa, plasma, human, use of polarization of fluorescence to measure the dissociation of (Freyssinet, J.-M., Lewis, B. A., Holbrook, J. J. & Shore, J. D.) 403–410

Coagulation Factor, *see* Clotting Factor

Coenzyme A, specificity of activation by acyl derivatives of thermophilic *Bacillus* sp. pyruvate carboxylase and properties of catalysis by the enzyme in the absence of activator (Libor, S. M., Sundaram, T. K. & Scrutton, M. C.) 543–558

Colchine-binding protein, *Dictyostelium discoideum*, characterization of (Cappuccinelli, P. & Hames, B. D.) 499–504

Collagen galactosyltransferase, chick-embryo, characterization of (Risteli, L.) 189–196

Collagenase, skin, human, purification, characterization and inhibition of (Woolley, D. E., Glanville, R. W., Roberts, D. R. & Evanson, J. M.) 265–276

Collagens, identification of the sites i[n] α-chains of, from various sources that bind guinea-pig and human serum anti-gelatin factor (cold-insoluble globulin) (Dessau, W., Adelmann, B. C., Timpl, R. & Martin, G. R.) 55–59

Copper ions, binding of, by a tripeptide corresponding to the copper(II)-binding site of human serum albumin (Iyer, K. S., Lau, S.-J., Laurie, S. H. & Sarkar, B.) 61–69
Copper ions, kinetic aspects of the binding of, to copper-free ox superoxide dismutase (Rigo, A., Viglino, P., Bonori, M., Cocco, D., Calabrese, L. & Rotilio, G.) 277–280
Coproporphyrinogen III, factors determining the sequence of oxidative decarboxylation of the 2- and 4-propionate substituents of, by rat liver coproporphyrinogen oxidase (Elder, G. H., Evans, J. O., Jackson, J. R. & Jackson, A. H.) 215–223
Coproporphyrinogen III, utilization of substrates other than, by rat liver coproporphyrinogen oxidase (Elder, G. H. & Evans, J. O.) 205–214
Coproporphyrinogen oxidase, liver, rat, factors determining the sequence of oxidative decarboxylation of the 2- and 4-propionate substituents of coproporphyrinogen III by (Elder, G. H., Evans, J. O., Jackson, J. R. & Jackson, A. H.) 215–223
Coproporphyrinogen oxidase, liver, rat, radiochemical method for assay of the activity of, and utilization of substrates other than coproporphyrinogen III by the enzyme (Elder, G. H. & Evans, J. O.) 205–214
Corneal stroma, ox, characterization of keratan sulphate-containing proteoglycans from (Axelsson, I. & Heinegård, D.) 517–530
Corpus luteum, ovary, cow, effects of guanosine triphosphate and other compounds on the activity of adenylate cyclase of (Young, J. L. & Stansfield, D. A.) 133–142
Cytochrome *c*, heart, horse, properties of, modified by attachment to a carbohydrate polymer (Silvestrini, M. C., Colosimo, A., Brunori, M. & Antonini, E.) 257–263
Cytochrome *c* oxidase, *Saccharomyces cerevisiae*, use of detergents for removal of lipids from, and reconstitution of enzyme activity by using cholate-mediated exchange (Virji, M. & Knowles, P. F.) 343–353
Cytosol, heart, pig, preparation of malate dehydrogenase from, and the binding of reduced nicotinamide–adenine dinucleotide by the enzyme (Lodola, A., Spragg, S. P. & Holbrook, J. J.) 577–588
Cytosol, liver, rat, physical properties of glucocorticoid-receptor protein from (Litwack, G., Cake, M. H., Filler, R. & Taylor, K.) 445–448
Cytosol, spleen, rat, partial purification and properties of acid nucleotidase from (Tjernshaugen, H.) 597–605

Deoxyribonucleic acid, mitochondrial, liver, rat, localization of the ribonucleotide sites in (Lonsdale, D. M. & Jones, I. G.) 79–85
Dictyostelium discoideum, characterization of the colchicine-binding protein from (Cappuccinelli, P. & Hames, B. D.) 499–504
2,4-Dinitrophenyl-binding myeloma protein (immunoglobulin A), binding of 2,4,6-trinitrophenyl derivatives to, from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Gettins, P., Jackson, R., Dwek, R. A. & Givol, D.) 179–188
Dodecyl sulphate, sodium, interaction of ox brain myelin basic protein with (Jones, A. J. S. & Rumsby, M. G.) 281–285
Dolichol, presence of, in *Saccharomyces cerevisiae* lipid diphosphate *N*-acetylglucosamine (Reuvers, F., Boer, P. & Hemming, F. W.) 505–508

Endoglucanase (cellulase), *Aspergillus niger*, substrate specificity and mode of action of (Hurst, P. L., Sullivan, P. A. & Shepherd, M. G.) 389–395
Enzymes, use of measurements of flux ratios for the determination of the order of addition of substrates and products in reactions catalysed by (Britton, H. G. & Dann, L. G.) 29–37
Escherichia coli, recognition of individual transfer ribonucleic acids from, by rat liver 1-adenine-specific transfer ribonucleic acid methyltransferase (Kraus, J.) 247–249
Escherichia coli (Crookes' strain), kinetic and structural properties of class-I fructose 1,6-bisphosphate aldolase from (Baldwin, S. A & Perham, R. N.) 643–652
Escherichia coli (Crookes' strain), purification and characterization of class-II fructose 1,6-bisphosphate aldolase from (Baldwin, S. A., Perham, R. N. & Stribling, D.) 633–641
Estradiol, *see* Oestradiol
Estrogens, *see* Oestrogens
Ethylenediaminetetra-acetic acid, demineralization of bone in organic solvents by alkylammonium salts of (Scott, J. E. & Kyffin, T. W.) 697–701
Euglena gracilis, fractionation of histones isolated from (Jardine, N. J. & Leaver, J. L.) 103–111
Eye, binding sites for retinoic acid in different tissues of, from various species (Wiggert, B., Bergsma, D. R., Helmsen, R. & Chader, G. J.) 87–94
Eye, ox, characterization of keratan sulphate-containing proteolgycans form corneal stroma of (Axelsson, I. & Heinegård, D.) 517–530

Factor XIII, plasma, human, equilibrium study of the binding of metal ions to (Lewis, B. A., Freyssinet, J.-M. & Holbrook, J. J.) 397–402
Factor XIIIa, plasma, human, use of polarization of fluorescence to measure the dissociation of (Freyssinet, J.-M., Lewis, B. A., Holbrook, J. J. & Shore, J. D.) 403–410
Fibrinogen, plasma, rat, purification of, and its constituent chains (van Ruijven-Vermeer, I. A. M. & Nieuwenhuizen, W.) 653–658
Fibroblasts, chick-embryo, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from, and other vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542
Fibroblasts, chick-embryo, young and senescent, content of actin in (Anderson, P. J.) 169–172
Flavin mononucleotide, covalently bound, participation of the iron–sulphur centres and, in the reaction catalysed by bacterium W3A1 trimethylamine dehydrogenase (Steenkamp, D. J., Singer, T. P. & Beinert, H.) 361–369
Fructose 1,6-bisphosphate aldolase, class-I, *Escherichia coli* (Crookes' strain), kinetic and structural properties of (Baldwin, S. A. & Perham, R. N.) 643–652
Fructose 1,6-bisphosphate aldolase, class-II, *Escherichia coli* (Crookes' strain), purification and characterization of (Baldwin, S. A., Perham, R. N. & Stribling, D.) 633–641

Gastric juice, human, use of diethylaminoethyl-cellulose for the preparation and purification of individual pepsins from (Roberts, N. B. & Taylor, W. H.) 607–615
Gastric mucosa, human, isolation and properties of a non-pepsin proteinase from (Roberts, N. B. & Taylor, W. H.) 617–624
Globulin, cold-insoluble (anti-gelatin factor), serum, guinea-pig and human, identification of the sites in α-chains of collagens from various sources that bind (Dessau, W., Adelmann, B. C., Timpl, R. & Martin, G. R.) 55–59
Glucocorticoid-receptor protein, cytosol, liver, rat, physical properties of (Litwack, G., Cake, M. H., Filler, R. & Taylor, K.) 445–448
Glutathione *S*-transferase B (ligandin), liver, rat, binding of porphyrins by (Tipping, E., Ketterer, B. & Koskelo, P.) 509–516
Glutathione *S*-transferases, activities of, in human lymphoid cells and fractionated peripheral leucocytes (Kaplowitz, N., Spina, C., Graham, M. & Kuhlenkamp, J.) 465–470
Glycoproteins, membrane, Thy-1, thymus-gland and brain, rat, molecular weights of, measured in the presence and absence of deoxycholate (Kuchel, P. W., Campbell, D. G., Barclay, A. N. & Williams, A. F.) 411–417
Growth hormone, *see* Somatotropin
p-Guanidino-L-phenylalanine, interaction of derivatives of, with thrombin and trypsin (Klausner, Y. S., Rigbi, M., Ticho, T., de Jong, P. J., Neginsky, E. J. & Rinott, Y.) 157–167
Guanosine triphosphate, effects of, and other compounds on the activity of cow ovary corpus-luteum adenylate cyclase (Young, J. L. & Stansfield, D. A.) 133–142

Haemoglobin, human, zinc ion-binding sites and affinity for oxygen of (Gilman, J. G. & Brewer, G. J.) 625–632
Heart, horse, properties of cytochrome *c* from, modified by attachment to a carbohydrate polymer (Silvestrini, M. C., Colosimo, A., Brunori, M. & Antonini, E.) 257–263
Heart, pig, inhibition by phosphorylation of additional sites on pyruvate dehydrogenase from, of re-activation of the enzyme by pyruvate dehydrogenase phosphate phosphatase (Sugden, P. H., Hutson, N. J., Kerbey, A. L. & Randle, P. J.) 433–435
Heart, pig, preparation of cytosol malate dehydrogenase from, and the binding of reduced nicotinamide–adenine dinucleotide by the enzyme (Lodola, A., Spragg, S. P. & Holbrook, J. J.) 577–588
HeLa cells, biosynthesis of a hypermodified nucleotide residue in *Saccharomyces cerevisiae* 17S ribosomal ribonucleic acid and in 18S ribonucleic acid of (Brand, R. C., Klootwijk, J., Planta, R. J. & Maden, B. E. H.) 71–77
HeLa cells, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from, and other vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542
Hen, *see* Chicken
Histones, *Euglena gracilis*, isolated, fractionation of (Jardine, N. J. & Leaver, J. L.) 103–111

Immunoglobulin A (myeloma protein), binding of 2,4,6-trinitrophenyl derivatives to, from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Gettins, P., Jackson, R., Dwek, R. A. & Givol, D.) 179–188
Instructions to authors, Policy of the Journal and 1–27
Iodination, comparison of the properties of guinea-pig low-density serum lipoproteins after, by two different methods (Weech, P. K., McTaggart, F. & Mills, G. L.) 687–695
Ions, gliding edge dislocations in p oteins as a possible mechanism for the active transport of, by biological membranes (Chandler, H. D., Woolf, C. J. & Hepburn, H. R.) 559–565
Iron–sulphur centres, magnetic coupling of the molybdenum and, in milk xanthine oxidase and in turkey liver and *Veillonella alcalescens* xanthine dehydrogenases (Lowe, D. J. & Bray, R. C.) 471–479
Iron–sulphur centres, participation of covalently bound flavin mononucleotide and, in the reaction catalysed by bacterium W3A1 trimethylamine dehydrogenase (Steenkamp, D. J., Singer, T. P. & Beinert, H.) 361–369

Keratan sulphate, characterization of ox corneal-stroma proteoglycans containing (Axelsson, I. & Heinegård, D.) 517–530
Kidney cells, baby-hamster, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from, and other vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542
Kidney cells, *Xenopus laevis*, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from, and other vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542
Kidney, rabbit, binding of ouabain to the membrane phospholipid-dependent sodium-plus-potassium ion-dependent adenosine triphosphatase from (Goodman, S. L. & Wheeler, K. P.) 313–320
Kidney, rabbit, simple and rapid method for the reversible removal of lipids from membrane sodium-plus-potassium ion-dependent adenosine triphosphatase from (Goodman, S. L., Isern de Caldentey, M. & Wheeler, K. P.) 305–311
Kidney, rat, rapid method for the assay of the activity of aminopropyltransferases from, and other tissues (Hibasami, H. & Pegg, A. E.) 709–712

β-Lactam ring, stereochemistry of the formation of, during the biosynthesis of cephalosporin by *Cephalosporium acremonium* (Huddleston, J. A., Abraham, E. P., Young, D. W., Morecombe, D. J. & Sen, P. K.) 705–707
β-Lactamase I, *Bacillus cereus*, reversible inhibitors of (Kiener, P. A. & Waley, S. G.) 197–204
β-Lactamase, *Staphylococcus aureus* PC1, preferential nitration by tetranitromethane of a specific tyrosine residue of (Bristow, A. F. & Virden, R.) 381–388
Lactation, inhibition of aggregation and characterization of mouse mammary-gland oestrogen-receptor protein during, in the presence of sodium bromide (Auricchio, F., Rotondi, A., Schiavone, E. & Bresciani, F.) 481–488

Lanostane, use of 32-functionalized derivatives of, in chemical and enzymic studies on the characterization of intermediates during the removal of the 14α-methyl group in the biosynthesis of cholesterol by cell-free preparations from (Akhtar, M., Alexander, K., Boar, R. B., McGhie, J. F. & Barton, D. H. R.) 449–463

L-cells, mouse, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from, and other vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542

Leghaemoglobin, anomalous migration of, on sodium dodecyl sulphate/polyacrylamide-gel electrophoresis (Lehtovaara, P.) 251–253

Leucocytes, peripheral, human, fractionated, activities of glutathione *S*-transferases in human lymphoid cells and (Kaplowitz, N., Spina, C., Graham, M. & Kuhlenkamp, J.) 465–470

Ligandin, liver, rat, binding of porphyrins by (Tipping, E., Ketterer, B. & Koskelo, P.) 509–516

Ligands, theoretical analysis of the use of zonal gel filtration in the detection and purification of complexes of proteins with (Nimmo, I. A. & Bauermeister, A.) 437–440

Lipid diphosphate *N*-acetylglucosamine, *Saccharomyces cerevisiae*, presence of dolichol in (Reuvers, F., Boer, P. & Hemming, F. W.) 505–508

Lipids, simple and rapid method for the reversible removal of, from rabbit kidney membrane sodium-plus-potassium ion-dependent adenosine triphosphatase (Goodman, S. L., Isern de Caldentey, M. & Wheeler, K. P.) 305–311

Lipids, use of detergents for removal of, from *Saccharomyces cerevisiae* cytochrome *c* oxidase and reconstitution of enzyme activity by phospholipids by using cholate-mediated exchange (Virji, M. & Knowles, P. F.) 343–353

Lipoproteins, serum, low-density, guinea-pig, comparison of the properties of, after iodination by two different methods (Weech, P. K., McTaggart, F. & Mills, G. L.) 687–695

Liver, dog, cat, rat and mouse, characteristics of isoenzymes of mitochondrial alanine–glyoxylate aminotransferase from (Noguchi, T., Okuno, E., Takada, Y., Minatogawa, Y., Okai, K. & Kido, R.) 113–122

Liver, rabbit, structural requirements in the inhibition by opioid substances of the reaction catalysed by microsomal morphine uridine diphosphate glucuronyltransferase from (Sanchez, E., del Villar, E. & Tephly, T. R.) 173–177

Liver, rat, binding of porphyrins by ligandin from (Tipping, E., Ketterer, B. & Koskelo, P.) 509–516

Liver, rat, factors determining the sequence of oxidative decarboxylation of the 2- and 4-propionate substituents of coproporphyrinogen III by coproporphyrinogen oxidase from (Elder, G. H., Evans, J. O., Jackson, J. R. & Jackson, A. H.) 215–223

Liver, rat, localization of the ribonucleotide sites in mitochondrial deoxyribonucleic acid from (Londsale, D. M. & Jones, I. G.) 79–85

Liver, rat, partial purification and characterization of a lysosomal proteinase from, catalysing the inactivation of pyruvate dehydrogenase complex (Lynen, A., Sedlaczek, E. & Wieland, O. H.) 321–328

Liver, rat, partial purification and properties of cytosol acid nucleotidases from, and spleen (Tjernshaugen, H.) 597–605

Liver, rat, physical properties of cytosol glucocorticoid-receptor protein from (Litwack, G., Cake, M. H., Filler, R. & Taylor, K.) 445–448

Liver, rat, purification and properties of nuclear adenosine 3′:5′-cyclic monophosphate-dependent protein kinase associated with ribonucleic acid polymerase I from (Hirsch, J. & Martelo, O. J.) 355–359

Liver, rat, purification of reduced nicotinamide–adenine dinucleotide (phosphate) dehydrogenase from, and its role in the vitamin K (3-methyl-3-phytyl-1,4-napthaquinone)-dependent carboxylation reaction involved in the biosynthesis of prothrombin (Wallin, R., Gebhardt, O. & Prydz, H.) 95–101

Liver, rat, radiochemical method for assay of the activity of coproporphyrinogen oxidase from, and utilization of substrates other than coproporphyrinogen III by the enzyme (Elder, G. H. & Evans, J. O.) 205–214

Liver, rat, rapid method for the assay of the activity of aminopropyltransferases from, and other tissues (Hibasami, H. & Pegg, A. E.) 709–712

Liver, rat, recognition of individual *Escherichia coli* transfer ribonucleic acids by 1-adenine-specific transfer ribonucleic acid methyltransferase from (Kraus, J.) 247–249

Liver, rat, use of 32-functionalized derivatives of lanostane in chemical and enzymic studies on the characterization of intermediates during the removal of the 14α-methyl group in the biosynthesis of cholesterol by cell-free preparations from (Akhtar, M., Alexander, K., Boar, R. B., McGhie, J. F. & Barton, D. H. R.) 449–463

Liver, sheep, pathway of the reaction catalysed by mitochondrial pyruvate carboxylase from (Easterbrook-Smith, S. B., Wallace, J. C. & Keech, D. B.) 225–228

Liver, turkey, magnetic coupling of the molybdenum and iron–sulphur centres in milk xanthine oxidase and in xanthine dehydrogenases from *Veillonella alcalescens* and (Lowe, D. J. & Bray, R. C.) 471–479

Lymphoid cells, human, activities of glutathione *S*-transferases in fractionated human peripheral leucocytes and (Kaplowitz, N., Spina, C., Graham, M. & Kuhlenkamp, J.) 465–470

Lysosomes, liver, rat, partial purification and characterization of a proteinase from, catalysing the inactivation of pyruvate dehydrogenase complex (Lynen, A., Sedlaczek, E. & Wieland, O. H.) 321–328

Malate dehydrogenase, cytosol, heart, pig, preparation of, and the binding of reduced nicotinamide–adenine dinucleotide by the enzyme (Lodola, A., Spragg, S. P. & Holbrook, J. J.) 577–588

Mammary gland, mouse, lactating, inhibition of aggregation and characterization of the oestrogen-receptor protein from, in the presence of sodium bromide (Auricchio, F., Rotondi, A., Schiavone, E. & Bresciani, F.) 481–488

Membrane, kidney, rabbit, binding of ouabain to the phospholipid-dependent sodium-plus-potassium ion-dependent adenosine triphosphatase of (Goodman, S. L. & Wheeler, K. P.) 313–320

Membrane, kidney, rabbit, simple and rapid method for the reversible removal of lipids from sodium-plus-potassium ion-dependent adenosine triphosphatase of (Goodman, S. L., Isern de Caldentey, M. & Wheeler, K. P.) 305–311

Membranes, biological, gliding edge dislocations in proteins as a possible mechanism for the active transport of ions by (Chandler, H. D., Woolf, C. J. & Hepburn, H. R.) 559–565

Membranes, thymus-gland and brain, rat, molecular weights of the Thy-1 glycoproteins from, measured in the presence and absence of deoxycholate (Kuchel, P. W., Campbell, D. G., Barclay, A. N. & Williams, A. F.) 411–417

Methanol dehydrogenases, dye-linked, partial purification of *Rhodopseudomonas acidophila* dye-linked alcohol dehydrogenase and comparison of its properties with those of, from other micro-organisms (Bamforth, C. W. & Quayle, J. R.) 677–686

14α-Methyl group, use of 32-functionalized derivatives of lanostane in chemical and enzymic studies on the characterization of intermediates during the removal of, in the biosynthesis of cholesterol by rat liver cell-free preparations (Akhtar, M., Alexander, K., Boar, R. B., McGhie, J. F. & Barton, D. H. R.) 449–463

3-Methyl-3-phytyl-1,4-naphthaquinone (vitamin K), purification of rat liver reduced nicotinamide–adenine dinucleotide (phosphate) dehydrogenase and its role in the carboxylation reaction dependent on, involved in the biosynthesis of prothrombin (Wallin, R., Gebhardt, O. & Prydz, H.) 95–101

Micrococcus lysodeikticus N.C.T.C. 2665, effect of temperature on the circular dichroism, fluorescence and enzymic activity of adenosine triphosphatase from (Ayala, J. A. & Nieto, M.) 371–380

Micrococcus sp. 2102, linkage of sugar phosphate polymer to peptidoglycan in cell walls of (Heptinstall, J., Coley, J., Ward, P. J., Archibald, A. R. & Baddiley, J.) 329–336

Microsomal fraction, liver, rabbit, structural requirements in the inhibition by opioid substances of the reaction catalysed by morphine uridine diphosphate glucuronyltransferase from (Sanchez, E., del Villar, E. & Tephly, T. R.) 173–177

Milk, human, isolation, composition and ordering of the peptides obtained by cyanogen bromide cleavage of casein from (Groves, M. L. & Greenberg, R.) 337–342

Milk, magnetic coupling of the molybdenum and iron–sulphur centres in turkey liver and *Veillonella alcalescens* xanthine dehydrogenases and in xanthine oxidase from (Lowe, D. J. & Bray, R. C.) 471–479

Mitochondria, liver, dog, cat, rat and mouse, characteristics of isoenzymes of alanine–glyoxylate aminotransferase from (Noguchi, T., Okuno, E., Takada, Y., Minatogawa, Y., Okai, K. & Kido, R.) 113–122

Mitochondria, liver, rat, localization of the ribonucleotide sites in deoxyribonucleic acid from (Lonsdale, D. M. & Jones, I. G.) 79–85

Mitochondria, liver, sheep, pathway of the reaction catalysed by pyruvate carboxylase from (Easterbrook-Smith, S. B., Wallace, J. C. & Keech, D. B.) 225–228

Molybdenum, magnetic coupling of the iron–sulphur centres and, in milk xanthine oxidase and in turkey liver and *Veillonella alcalescens* xanthine dehydrogenases (Lowe, D. J. & Bray, R. C.) 471–479

Mrophine uridine diphosphate glucuronyltransferase microsomal, liver, rabbit, structural requirements in the inhibition by opioid substances of the reaction catalysed by (Sanchez, E., del Villar, E. & Tephly, T. R.) 173–177

Mucosa, gastric, human, isolation and properties of a non-pepsin proteinase from (Roberts, B. N. & Taylor, W. H.) 617–624

Muscle, chick-embryo, isolation and separation of α- and β-tubulins from, and other tissues (El-Hamalawi, A.-R. A.) 717–719

Muscle, skeletal, breast and leg, chicken, comparison of troponin T and troponin I from (Wilkinson, J. M.) 229–238

Muscle, skeletal, rabbit, aggregational state of deoxycholate-solubilized sarcoplasmic-reticulum calcium-plus-magnesium ion-dependent adenosine triphosphatase from, in an enzymically active form (Jørgensen, K. E., Lind, K. E., Røigaard-Petersen, H. & Møller, J. V.) 489–498

Muscle, skeletal, rabbit, gliding edge dislocations in proteins as a possible mechanism for the active transport of ions by sarcoplasmic-reticulum membranes from, and by other biological membranes (Chandler, H. D., Woolf, C. J. & Hepburn, H. R.) 559–565

Muscle, skeletal, rabbit, kinetics and mechanism of action of pyruvate kinase from (Dann, L. G. & Britton, H. G.) 39–54

Myelin, brain, ox, cat, rabbit and rat, proteolytic and peroxidatic reactions of commercial preparations of horseradish peroxidase with basic protein of (Cammer, W., Bieler, L. Z. & Norton, W. T.) 567–577

Myelin, brain, ox, interaction of sodium dodecyl sulphate with basic protein from (Jones, A. J. S. & Rumsby, M. G.) 281–285

Myeloma protein (immunoglobulin A), binding of 2,4,6-trinitrophenyl derivatives to, from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Gettins, P., Jackson, R., Dwek, R. A. & Givol, D.) 179–188

Nasal septum, ox, characterization of proteinase inhibitors of cartilage of (Roughley, P. J., Murphy, G. & Barrett, A. J.) 721–724

Neurospora crassa, evidence from peptide 'maps' for the identity of subunits of the *arom* multienzyme complex from (Lumsden, J. & Coggins, J. R.) 441–444

Nicotinamide–adenine dinucleotide (phosphate) (reduced) dehydrogenase, liver, rat, purification of, and its role in the vitamin K (3-methyl-3-phytyl-1,4-naphthaquinone)-dependent carboxylation reaction involved in the biosynthesis of prothrombin (Wallin, R., Gebhardt, O. & Prydz, H.) 95–101

Nicotinamide–adenine dinucleotide, reduced, binding of, by pig heart cytosol malate dehydrogenase (Lodola, A., Spragg, S. P. & Holbrook, J. J.) 577–588

Nuclei, *Euglena gracilis*, fractionation of histones isolated from (Jardine, N. J. & Leaver, J. L.) 103–111

Nuclei, liver, rat, purification and properties of adenosine 3′:5′-cyclic monophosphate-dependent protein kinase associated with ribonucleic acid polymerase I from (Hirsch, J. & Martelo, O. J.) 355–359

Nucleoside diphosphatase, pea-seedling, partial purification and properties of, and uridine diphosphate glucose–sterol glucosyltransferase (Staver, M. J., Glick, K. & Baisted, D. J.) 297–303
Nucleotidase, acid, cytosol, spleen, rat, partial purification and properties of (Thernshaugen, H.) 597–605
Nucleotide sequences, methylated, extensive homologies between, of ribosomal ribonucleic acids from various vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542

Oestradiol-17β-receptor protein, mammary-gland, mouse, lactating, inhibition of aggregation and characterization of, in the presence of sodium bromide (Auricchio, F., Rotondi, A., Schiavone, E. & Bresciani, F.) 481–488
Oestrogen-receptor protein, mammary-gland, mouse, lactating, inhibition of aggregation and characterization of, in the presence of sodium bromide (Auricchio, F., Rotondi, A., Schiavone, E. & Bresciani, F.) 481–488
Osteoarthritis, changes in the composition of dog articular-cartilage proteoglycans during aging and (Inerot, S., Heinegård, D., Audell, L. & Olsson, S.-E.) 143–156
Ouabain, binding of, to rabbit kidney membrane phospholipid-dependent sodium-plus-potassium ion-dependent adenosine triphosphatase (Goodman, S. L. & Wheeler, K. P.) 313–320
Ovary, cow, effects of guanosine triphosphate and other compounds on the activity of adenylate cyclase of corpus luteum of (Young, J. L. & Stansfield, D. A.) 133–142
Oxygen, zinc ion-binding sites and affinity for, of human haemoglobin (Gilman, J. G. & Brewer, G. J.) 625–632

Pea (*Pisum sativum*) seedlings, partial purification and properties of uridine diphosphate glucose–sterol glucosyltransferase and nucleoside diphosphatase from (Staver, M. J., Glick, K. & Baisted, D. J.) 297–303
Penicillinase (β-lactamase), *Staphylococcus aureus* PC1, preferential nitration by tetranitromethane of a specific tyrosine residue of (Bristow, A. F. & Virden, R.) 381–388
Penicillinase (β-lactamase I), *Bacillus cereus*, reversible inhibitors of (Kiener, P. A. & Waley, S. G.) 197–204
Pepsins, gastric-juice, human, individual, use of diethylaminoethyl-cellulose for the preparation and purification of (Roberts, N. B. & Taylor, W. H.) 607–615
Peptides, isolation, composition and ordering of, obtained by cyanogen bromide cleavage of human casein (Groves, M. L. & Greenberg, R.) 337–342
Peptidoglycan, linkage of sugar phosphate polymer to, in *Micrococcus* sp. 2102 cell wall (Heptinstall, J., Coley, J., Ward, P. J., Archibald, A. R. & Baddiley, J.) 329–336
Peroxidase, horseradish, proteolytic and peroxidatic reactions of commercial preparations of, with ox, cat, rabbit and rat brain myelin basic protein (Cammer, W., Bieler, L. Z. & Norton, W. T.) 567–577
Phospholipids, simple and rapid method for the reversible removal of, from rabbit kidney membrane sodium-plus-potassium ion-dependent adenosine triphosphatase (Goodman, S. L., Isern de Caldentey, M. & Wheeler, K. P.) 305–311
Phospholipids, use of detergents for removal of lipids from *Saccharomyces cerevisiae* cytochrome *c* oxidase and reconstitution of enzyme activity by, by using cholate-mediated exchange (Virji, M. & Knowles, P. F.) 343–353
Pigment epithelium, binding sites for retinoic acid in, and other ocular tissues from various species (Wiggert, B., Bergsma, D. R., Helmsen, R. & Chader, G. J.) 87–94
Pisum sativum, *see* Pea
Pituitary gland, anterior, rat, preparation of ^{14}C-labelled somatotropin by biosynthesis by, *in vitro* (Wallis, M., Davies, R. V. & Daniels, M.) 669–676
Plasma-cell tumour MOPC 315 cells, mouse, binding of 2,4,6-trinitrophenyl derivatives to myeloma protein (immunoglobulin A) from (Dower, S. K., Gettins, P., Jackson, R., Dwek, R. A. & Givol, D.) 179–188
Plasma, human, equilibrium study of the binding of metal ions to Factor XIII from (Lewis, B. A., Freyssinet, J.-M. & Holbrook, J. J.) 397–402
Plasma, human, use of polarization of fluorescence to measure the dissociation of Factor XIIIa from (Freyssinet, J.-M., Lewis, B. A., Holbrook, J. J. & Shore, J. D.) 403–410
Policy of the Journal and instructions to authors 1–27
Plasma, rabbit, isolation and partial characterization of α_1-anti-trypsin from (Koj, A., Hatton, M. W. C., Wong, K.-L. & Regoeczi, E.) 589–596
Plasma, rat, purification of fibrinogen and its constituent chains from (van Ruijven-Vermeer, I. A. M. & Nieuwenhuizen, W.) 653–658
Poly-(*N*-acetylglucosamine 1-phosphate), linkage of, to peptidoglycan in *Micrococcus* sp. 2102 cell wall (Heptinstall, J., Coley, J., Ward, P. J., Archibald, A. R. & Baddiley, J.) 329–336
Polysaccharide, properties of horse heart cytochrome *c* modified by attachment to (Silvestrini, M. C., Colosimo, A., Brunori, M. & Antonini, E.) 257–263
Porphyrins, binding of, by rat liver ligandin (Tipping, E., Ketterer, B. & Koskelo, P.) 509–516
Prostate gland, ventral, rat, rapid method for the assay of the activity of aminopropyltransferases from, and other tissues (Hibasami, H. & Pegg, A. E.) 709–712
Protease, *see* Proteinase
Protein, basic, myelin, brain, ox, cat, rabbit and rat, proteolytic and peroxidatic reactions of commercial preparations of horseradish peroxidase with (Cammer, W., Bieler, L. Z. & Norton, W. T.) 567–577
Protein, basic, myelin, brain, ox, interaction of sodium dodecyl sulphate with (Jones, A. J. S. & Rumsby, M. G.) 281–285
Protein, colchicine-binding, *Dictyostelium discoideum*, characterization of (Cappuccinelli, P. & Hames, B. D.) 499–504
Protein, glucocorticoid-receptor, cytosol, liver, rat, physical properties of (Litwack, G., Cake, M. H., Filler, R. & Taylor, K.) 445–448
Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, nuclear, liver, rat, purification and properties of, associated with ribonucleic acid polymerase I (Hirsch, J. & Martelo, O. J.) 355–359
Protein, myeloma (immunoglobulin A), binding of 2,4,6-trinitrophenyl derivatives to, from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Gettins, P., Jackson, R., Dwek, R. A. & Givol, D.) 179–188

Protein, oestrogen-receptor, mammary-gland, mouse, lactating, inhibition of aggregation and characterization of, in the presence of sodium bromide (Auricchio, F., Rotondi, A., Schiavone, E. & Bresciani, F.) 481–488
Proteinase inhibitors, cartilage, nasal-septum, ox, characterization of (Roughley, P. J., Murphy, G. & Barrett, A. J.) 721–724
Proteinase, lysosomal, liver, rat, partial purification and characterization of, catalysing the inactivation of pyruvate dehydrogenase complex (Lynen, A., Sedlaczek, E. & Wieland, O. H.) 321–328
Proteinase, non-pepsin, gastric-mucosa, human, isolation and properties of (Roberts, N. B. & Taylor, W. H.) 617–624
Proteinases, skin, rabbit, two, characterization of (Jasani, B., Jasani, M. K. & Talbot, M. D.) 287–295
Protein–polysaccharides, *see* Proteoglycans
Proteins, gliding edge dislocations in, as a possible mechanism for the active transport of ions by biological membranes (Chandler, H. D., Woolf, C. J. & Hepburn, H. R.) 559–565
Proteins, theoretical analysis of the use of zonal gel filtration in the detection and purification of complexes of, with ligands (Nimmo, I. A. & Bauermeister, A.) 437–440
Proteoglycans, articular-cartilage, dog, changes in the composition of, during aging and osteoarthritis (Inerot, S., Heinegård, D., Audell, L. & Olsson, S.-E.) 143–156
Proteoglycans, articular-cartilage, human, isolation of (Bayliss, M. T. & Ali, S. Y.) 123–132
Proteoglycans, keratan sulphate-containing, corneal-stroma, ox, characterization of (Axelsson, I. & Heinegård, D.) 517–530
Proteoglycans, preservation of, in bone matrix during demineralization in organic solvents by alkylammonium salts of ethylenediaminetetra-acetic acid (Scott, J. E. & Kyffin, T. W.) 697–701
Prothrombin, purification of rat liver reduced nicotinamide–adenine dinucleotide (phosphate) dehydrogenase and its role in the vitamin K (3-methyl-3-phytyl-1,4-naphthaquinone)-dependent carboxylation reaction involved in the biosynthesis of (Wallin, R., Gebhardt, O. & Prydz, H.) 95–101
Pseudomonas C12B, purification and properties of S1 secondary alklysulphatase from (Bartholomew, B., Dodgson, K. S. & Gorham, S. D.) 659–667
Pseudomonas MA-1, determination of the amino acid sequence around the pyridoxal-binding site of pyridoxamine–pyruvate aminotransferase from (Hodsdon, J., Kolb, H., Snell, E. E. & Cole, R. D.) 429–432
Pyridoxal, determination of the amino acid sequence around the binding site for, of *Pseudomonas* MA-1 pyridoxamine–pyruvate aminotransferase (Hodsdon, J., Kolb, H., Snell, E. E. & Cole, R. D.) 429–432
Pyridoxamine–pyruvate aminotransferase, *Pseudomonas* MA-1, determination of the amino acid sequence around the pyridoxal-binding site of (Hodsdon, J., Kolb, H., Snell, E. E. & Cole, R. D.) 429–432
Pyruvate carboxylase, *Bacillus* sp., thermophilic, specificity of activation of, by acyl-coenzyme A and properties of catalysis by the enzyme in the absence of activator (Libor, S. M., Sundaram, T. K. & Scrutton, M. C.) 543–558
Pyruvate carboxylase, mitochondrial, liver, sheep, pathway of the reaction catalysed by (Easterbrook-Smith, S. B., Wallace, J. C. & Keech, D. B.) 225–228
Pyruvate dehydrogenase complex, partial purification and characterization of a rat liver lysosomal proteinase catalysing the inactivation of (Lynen, A., Sedlaczek, E. & Wieland, O. H.) 321–328
Pyruvate dehydrogenase, heart, pig, inhibition by phosphorylation of additional sites on, of re-activation of the enzyme by pyruvate dehydrogenase phosphate phosphatase (Sugden, P. H., Hutson, N. J., Kerbey, A. L. & Randle, P. J.) 433–435
Pyruvate kinase, skeletal-muscle, rabbit, kinetics and mechanism of action of (Dann, L. G. & Britton, H. G.) 39–54

Rennin (chymosin), demonstration of the activity of, in the stomach of the newborn pig (Foltmann, B., Lønblad, P. & Axelsen, N. H.) 425–427
Reticulum, sarcoplasmic, skeletal-muscle, rabbit, aggregational state of deoxycholate-solubilized calcium-plus-magnesium ion-dependent adenosine triphosphatase from, in an enzymically active form (Jørgensen, K. E., Lind, K. E., Røigaard-Petersen, H. & Møller, J. V.) 489–498
Reticulum, sarcoplasmic, skeletal-muscle, rabbit, gliding edge dislocations in proteins as a possible mechanism for the active transport of ions by, and by other biological membranes (Chandler, H. D., Woolf, C. J. & Hepburn, H. R.) 559–565
Retina, binding sites for retinoic acid in, and other ocular tissues from various species (Wiggert, B., Bergsma, D. R., Helmsen, R. & Chader, G. J.) 87–94
Retinoic acid, binding sites for, in different ocular tissues from various species (Wiggert, B., Bergsma, D. R., Helmsen, R. & Chader, G. J.) 87–94
Rhodopseudomonas acidophila, partial purification of dye-linked alcohol dehydrogenase from, and comparison of its properties with those of dye-linked methanol dehydrogenases from other micro-organisms (Bamforth, C. W. & Quayle, J. R.) 677–686
Ribonucleic acid, kinetics and template-dependency of the biosynthesis of, by bacterial ribonucleic acid polymerases (Panayotatos, N. & Kèzdy, F. J.) 421–423
Ribonucleic acid polymerase I, nuclear, liver, rat, purification and properties of adenosine 3′:5′-cyclic monophosphate-dependent protein kinase associated with (Hirsch, J. & Martelo, O. J.) 355–359
Ribonucleic acid polymerases, bacterial, kinetics and template-dependency of the biosynthesis of ribonucleic acid by (Panayotatos, N. & Kèzdy, F. J.) 421–423
Ribonucleic acid, ribosomal, 17S, *Saccharomyces cerevisiae*, biosynthesis of a hypermodified nucleotide residue in HeLa-cell 18S ribosomal ribonucleic acid and in (Brand, R. C., Klootwijk, J., Planta, R. J. & Maden, B. E. H.) 71–77
Ribonucleic acid, ribosomal, 18S, HeLa-cell, biosynthesis of a hypermodified nucleotide residue in *Saccharomyces cerevisiae* 17S ribosomal ribonucleic acid and in (Brand, R. C., Klootwijk, J., Planta, R. J. & Maden, B. E. H.) 71–77
Ribonucleic acid (transfer) methyltransferase, liver, rat, recognition of individual *Escherichia coli* transfer ribonucleic acids by (Kraus, J.) 247–249

Ribonucleic acids, ribosomal, extensive homologies between the methylated nucleotide sequences of, from various vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542
Ribonucleic acids, transfer, *Escherichia coli*, recognition of, by rat liver 1-adenine-specific transfer ribonucleic acid methyltransferase (Kraus, J.) 247–249
Ribonucleotide sites, localization of, in rat liver mitochondrial deoxyribonucleic acid (Lonsdale, D. M. & Jones, I. G.) 79–85
Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal

Saccharomyces cerevisiae, biosynthesis of a hypermodified nucleotide residue in HeLa-cell 18S ribosomal ribonucleic acid and in 17S ribosomal ribonucleic acid of (Brand, R. C., Klootwijk, J., Planta, R. J. & Maden, B. E. H.) 71–77
Saccharomyces cerevisiae, presence of dolichol in a lipid diphosphate *N*-acetylglucosamine from (Reuvers, F., Boer, P. & Hemming, F. W.) 505–508
Saccharomyces cerevisiae, use of detergents for removal of lipids from cytochrome *c* oxidase from, and reconstitution of enzyme activity by using cholate-mediated exchange (Virji, M. & Knowles, P. F.) 343–353
Sarcoplasmic reticulum, *see* Reticulum, sarcoplasmic
Seedlings, pea, partial purification and properties of uridine diphosphate glucose–sterol glucosyltransferase and nucleoside diphosphatase from (Staver, M. J., Glick, K. & Baisted, D. J.) 297–303
Serum albumin, *see* Albumin, serum
Serum, guinea-pig and human, identification of the sites in α-chains of collagens from various sources that bind anti-gelatin factor (cold-insoluble globulin) from (Dessau, W., Adelmann, B. C., Timpl, R. & Martin, G. R.) 55–59
Serum lipoproteins, *see* Lipoproteins, serum
Skeletal muscle, *see* Muscle, skeletal
Skin, chick-embryo, isolation and separation of α- and β-tubulins from, and other tissues (El-Hamalawi, A.-R. A.) 717–719
Skin, human, purification, characterization and inhibition of collagenase from (Woolley, D. E., Glanville, R. W., Roberts, D. R. & Evanson, J. M.) 265–276
Skin, rabbit, characterization of two proteinases found in homografts of (Jasani, B., Jasani, M. K. & Talbot, M. D.) 287–295
Slime mould (*Dictyostelium discoideum*), characterization of the colchicine-binding protein from (Cappuccinelli, P. & Hames, B. D.) 499–504
Sodium dodecyl sulphate, interaction of ox brain myelin basic protein with (Jones, A. J. S. & Rumsby, M. G.) 281–285
Somatotropin, rat, ^{14}C-labelled, preparation of, by biosynthesis by rat anterior pituitary gland *in vitro* (Wallis, M., Davies, R. V. & Daniels, M.) 669–676
Spleen, rat, partial purification and properties of cytosol acid nucleotidase from (Tjernshaugen, H.) 597–605
Staphylococcus aureus PC1, preferential nitration by tetranitromethane of a specific tyrosine residue in β-lactamase from (Bristow, A. F. & Virden, R.) 381–388
Stomach, human, isolation and properties of a non-pepsin proteinase from the mucosa of (Roberts, N. B. & Taylor, W. H.) 617–624
Stomach, human, use of diethylaminoethyl-cellulose for the preparation and purification of individual pepsins from gastric juice from (Roberts, N. B. & Taylor, W. H.) 607–615
Stomach, pig, newborn, demonstration of the activity of rennin in (Foltmann, B., Lønblad, P. & Axelsen, N. H.) 425–427
Sugar phosphate polymer, linkage of, to peptidoglycan in *Micrococcus* sp. 2102 cell wall (Heptinstall, J., Coley, J., Ward, P. J., Archibald, A. R. & Baddiley, J.) 329–336
Supernatant fraction, *see* Cytosol
Superoxide dismutase, ox, copper-free, kinetic aspects of the binding of copper ions to (Rigo, A., Viglino, P., Bonori, M., Cocco, D., Calabrese, L. & Rotilio, G.) 277–280

Temperature, effect of, on the circular dichroism, fluorescence and enzymic activity of *Micrococcus lysodeikticus* N.C.T.C. 2665 adenosine triphosphatase (Ayala, J. A. & Nieto, M.) 371–380
Tetranitromethane, preferential nitration by, of a specific tyrosine residue of *Staphylococcus aureus* PC1 β-lactamase (Bristow, A. F. & Virden, R.) 381–388
Thrombin, interaction of α-*N*-(toluene-*p*-sulphonyl)-*p*-guanidino-L-phenylalanine methyl ester with trypsin and (Klausner, Y. S., Rigbi, M., Ticho, T., de Jong, P. J., Neginsky, E. J. & Rinott, Y.) 157–167
Thymus gland, rat, molecular weights of the Thy-1 membrane glycoproteins from, and brain measured in the presence and absence of deoxycholate (Kuchel, P. W., Campbell, D. G., Barclay, A. N. & Williams, A. F.) 411–417
α-*N*-(Toluene-*p*-sulphonyl)-*p*-guanidino-L-phenylalanine methyl ester, interaction of, with thrombin and trypsin (Klausner, Y. S., Rigbi, M., Tichi, T., de Jong, P. J., Neginsky, E. J. & Rinott, Y.) 157–167
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Trichoderma koningii, affinity chromatography of the components of cellulase system from (Halliwell, G. & Griffin, M.) 713–715
Trimethylamine dehydrogenase, bacterium W3A1, participation of the iron–sulphur centres and of covalently bound flavin mononucleotide in the reaction catalysed by (Steenkamp, D. J., Singer, T. P. & Beinert, H.) 361–369
2,4,6-Trinitrophenyl derivatives, binding of, to myeloma protein (immunoglobulin A) from mouse plasma-cell tumour MOPC 315 cells (Dower, S. K., Gettins, P., Jackson, R., Dwek, R. A. & Givol, D.) 179–188
Troponin I, skeletal-muscle, breast and leg, chicken, comparison of, and troponin T (Wilkinson, J. M.) 229–238
Troponin T, skeletal-muscle, breast and leg, chicken, comparison of, and troponin I (Wilkinson, J. M.) 229–238
Trypsin inhibitor (α_1-anti-trypsin), plasma, rabbit, isolation and partial characterization of (Koj, A., Hatton, M. W. C., Wong, K.-L. & Regoeczi, E.) 589–596
Trypsin, interaction of α-*N*-(toluene-*p*-sulphonyl)-*p*-guanidino-L-phenylalanine methyl ester with thrombin and (Klausner, Y. S., Rigbi, M., Ticho, T., de Jong, P. J., Neginsky, E. J. & Rinott, Y.) 157–167

Tubulin (colchicine-binding protein), *Dictyostelium discoideum*, characterization of (Cappuccinelli, P. & Hames, B. D.) 499–504
α- and β-Tubulins, brain, muscle and skin, chick-embryo, isolation and separation of (El-Hamalawi, A.-R. A.) 717–719
Tyrosine residue, specific, preferential nitration by tetranitromethane of, of *Staphyloccocus aureus* PC1 β-lactamase (Bristow, A. F. & Virden, R.) 381–388

Uridine diphosphate glucose–sterol glucosyltransferase, pea-seedling, partial purification and properties of, and nucleoside diphosphatase (Staver, M. J., Glick, K. & Baisted, D. J.) 297–303
Uridine diphosphate glucuronyltransferase, microsomal, liver, rabbit, structural requirements in the inhibition by opioid substances of the activity of, towards morphine (Sanchez, E., del Villar, E. & Tephly, T. R.) 173–177

Veillonella alcalescens, magnetic coupling of the molybdenum and iron–sulphur centres in milk xanthine oxidase and in xanthine dehydrogenases from turkey liver and (Lowe, D. J. & Bray, R. C.) 471–479
Vitamin A receptors, binding of retinoic acid by, in different ocular tissues from various species (Wiggert, B., Bergsma, D. R., Helmsen, R. & Chader, G. J.) 87–94
Vitamin K (3-methyl-3-phytyl-1,4-naphthaquinone), purification of rat liver reduced nicotinamide–adenine dinucleotide (phosphate) dehydrogenase and its role in the carboxylation reaction dependent on, involved in the biosynthesis of prothrombin (Wallin, R., Gebhardt, O. & Prydz, H.) 95–101

Wall, cell, *see* Cell wall

Xanthine dehydrogenases, *Veillonella alcalescens* and turkey liver, magnetic coupling of the molybdenum and iron–sulphur centres of milk xanthine oxidase and (Lowe, D. J. & Bray, R. C.) 471–479
Xanthine oxidase, milk, magnetic coupling of the molybdenum and iron–sulphur centres of turkey liver and *Veillonella alcalescens* xanthine dehydrogenases and (Lowe, D. J. & Bray, R. C.) 471–479
Xenopus laevis, extensive homologies between the methylated nucleotide sequences of ribosomal ribonucleic acids from kidney cells from, and other vertebrate cells (Khan, M. S. N., Salim, M. & Maden, B. E. H.) 531–542

Yeast, baker's (*Saccharomyces cerevisiae*), presence of dolichol in a lipid diphosphate *N*-acetylglucosamine from (Reuvers, F., Boer, P. & Hemming, F. W.) 505–508
Yeast, baker's (*Saccharomyces cerevisiae*), use of detergents for removal of lipids from cytochrome *c* oxidase from, and reconstitution of enzyme activity by phospholipids by using cholate-mediated exchange (Virji, M. & Knowles, P. F.) 343–353
Yeast (*Saccharomyces cerevisiae*), biosynthesis of a hypermodified nucleotide residue in HeLa-cell 18S ribosomal ribonucleic acid and in 17S ribonucleic acid of (Brand, R. C., Klootwijk, J., Planta, R. J. & Maden, B. E. H.) 71–77

Zinc ions, binding sites for, and affinity for oxygen of human haemoglobin (Gilman, J. G. & Brewer, G. J.) 625–632